

Table 6. Examples of preferred embodiments of the methods for modifying glycosylation in a eukaryotic microorganism, e.g. *Pichia pastoris*

<i>Desired Structure</i>	<i>Suitable Catalytic Activities</i>	<i>Suitable Sources of Localization Sequences</i>	<i>Suitable Gene Deletions</i>	<i>Suitable Transporters and/or Phosphatases</i>
Man <sub>5</sub> GlcNAc <sub>2</sub>	α-1,2-mannosidase (murine, human, <i>Bacillus</i> sp., <i>A. nidulans</i> )	Mns1 (N-terminus, <i>S. cerevisiae</i> ) Och1 (N-terminus, <i>S. cerevisiae</i> , <i>P. pastoris</i> ) Ktr1 Mnn9 Mnt1 ( <i>S. cerevisiae</i> ) KDEL (SEQ ID NO:6), HDEL (SEQ ID NO:5) (C-terminus)	OCH1 MNN4 MNN6	none
GlcNAcMan <sub>5</sub> GlcNAc <sub>2</sub>	GlcNAc Transferase I, (human, murine, rat etc.)	Och1 (N-terminus, <i>S. cerevisiae</i> , <i>P. pastoris</i> ) KTR1 (N-terminus) KDEL (SEQ ID NO:6), HDEL (SEQ ID NO:5) (C-terminus) Mnn1 (N-terminus, <i>S. cerevisiae</i> ) Mnt1 (N-terminus, <i>S. cerevisiae</i> ) GDPase (N-terminus, <i>S. cerevisiae</i> )	OCH1 MNN4 MNN6	UDP-GlcNAc transporter (human, murine, <i>K. lactis</i> ) UD Pase (human)
GlcNAcMan <sub>3</sub> GlcNAc	mannosidase II	Ktr1 Mnn1 (N-terminus,	OCH1	UDP-GlcNAc

Ac <sub>2</sub>		<i>S. cerevisiae</i> )	MNN4	transporter)
-----------------	--	------------------------	------	--------------